

Quantum Reference Frames and Quantum Transformations.

M. Toller

Dipartimento di Fisica dell'Università, Trento
I.N.F.N. gruppo collegato di Trento, Italia

February 7, 2008

Abstract

A quantum frame is defined by a material object subject to the laws of quantum mechanics. The present paper studies the relations between quantum frames, which in the classical case are described by elements of the Poincaré group. The possibility of using a suitable quantum group is examined, but some arguments are given which show that a different mathematical structure is necessary. Some simple examples in lower dimensional spacetimes are treated. They indicate the necessity of taking into account some “internal” degrees of freedom of the quantum frames, that can be disregarded in a classical treatment.

1 Introduction.

It has been stressed by several authors [1, 2, 3, 4] that, from the physical point of view, a frame of reference is defined by a material object of the same nature as the objects that form the system under investigation and the measuring instruments. Then, in principle, one should take into account:

1. The gravitational field generated by this object.
2. Its quantum properties that do not permit an exact determination of its position and of its velocity.

The gravitational field can be disregarded if the mass of the object that defines the reference frame is sufficiently small and the quantum effects are not important if the mass is sufficiently large. If, in a given physical situation, one can choose a mass that satisfies both these conditions, the material nature of the reference frames becomes irrelevant. Otherwise, we fall into the domain of quantum gravity and the usual geometric concepts are not valid any more [5, 6, 7, 8].

In the present work, as in ref. [2], we disregard the gravitational interaction and we consider quantum reference frames with finite mass M . Of course, the physical phenomena can be described completely by considering the limit $M \rightarrow \infty$. However, we think that a good understanding of this problem for finite M is preliminary to a treatment taking into account the effects of quantum gravity.

The relations between classical reference frames are described by elements of the Poincaré or of the Galilei group $\mathcal{G}$. Our aim is to find a mathematical structure that describes the relations between quantum frames. We start from the remark that in the absence of a classical reference frame the observable quantities, which concern both the objects that define a quantum reference frame and the objects described by the theory, have to be invariant under the action of the group $\mathcal{G}$.

Various authors [9, 10, 11, 12] have suggested that the quantum aspects of spacetime can be described by a quantum group [13, 14] obtained from a deformation of the commutative Hopf algebra of the functions defined on the group $\mathcal{G}$. From the point of view of the quantum frames, even in the absence of gravitation, there are two problems:

1. A quantum group cannot describe the “internal” degrees of freedom of the quantum frames and we shall see that they play an essential role.
2. If we have three quantum frames F_1 , F_2 and F_3 , the observables which describe the relation between F_1 and F_2 cannot be compatible (in the sense of quantum theory) with the observables which describe the relations between F_2 and F_3 . This fact cannot be taken into account by a quantum group.

The second problem is an aspect of the “paradox of quantum frames” discussed in ref. [2].

In Section 2 we introduce a general mathematical structure which describes the relations between quantum frames. In Section 3 we consider the classical (non quantum) case and we discuss the connection with the formalism based on Hopf algebras. In Sections 4, 5 and 6 we study some simple examples in zero and one space dimensions, both relativistic and non relativistic. The role of the “internal” degrees of freedom of the quantum frames is clarified by these examples.

It has been suggested in ref. [8] that in a quantum description of spacetime it is necessary to introduce some new degrees of freedom, besides the spacetime coordinates. We suggest that they can be interpreted as “internal” degrees of freedom of the quantum frames.

2 Relations between Quantum Frames.

We consider in a flat spacetime a system composed of n objects $F_0, \dots, F_{n-1}$. Some of them define reference frames and the others are the objects described by the theory. In order to simplify the problem, we assume that these objects do not interact mutually, but only with the measuring instruments. This assumption is justified only if we can disregard gravitation, a long range interaction which involves in an universal way all the kinds of physical objects.

In a quantum treatment, the states of the system are described by the Hilbert space

$$\mathcal{H}^{(n)} = \mathcal{H}_0 \otimes \mathcal{H}_1 \otimes \dots \otimes \mathcal{H}_{n-1}. \quad (1)$$

We assume that the bounded observables are described by Hermitean operators belonging to the algebra¹ $\mathcal{A}^{(n)} = \mathcal{L}(\mathcal{H}^{(n)})$ of the bounded operators in $\mathcal{H}^{(n)}$. The observables which concern the object F_k are described by the algebra $\mathcal{A}_k = \mathcal{L}(\mathcal{H}_k)$ and the observables which concern the pair of objects $\{F_i, F_k\}$ are described by the algebra $\mathcal{A}_{ik} = \mathcal{L}(\mathcal{H}_i \otimes \mathcal{H}_k)$. We define the injective homomorphisms $j_k^{(n)} : \mathcal{A}_k \rightarrow \mathcal{A}^{(n)}$ and $j_{ik}^{(n)} : \mathcal{A}_{ik} \rightarrow \mathcal{A}^{(n)}$ by means of the formulas

$$j_k^{(n)}(A_k) = B_0 \otimes \dots \otimes B_{n-1}, \quad B_k = A_k \in \mathcal{A}_k, \quad B_j = 1 \quad \text{for } j \neq k, \quad (2)$$

¹In a more refined treatment, one should consider a smaller C^* algebra $\mathcal{A}^{(n)} \subset \mathcal{L}(\mathcal{H}^{(n)})$, in order to take into account the possible occurrence of superselection rules and the topological properties of the system [15].

$$\begin{aligned}
j_{ik}^{(n)}(A_i \otimes A_k) &= B_0 \otimes \cdots \otimes B_{n-1}, & B_i &= A_i \in \mathcal{A}_i, & B_k &= A_k \in \mathcal{A}_k, \\
B_j &= 1 \quad \text{for } j \neq i, \quad j \neq k.
\end{aligned} \tag{3}$$

These homomorphisms formalize the concept of subsystem, which is essential for the following considerations.

We adopt the Heisenberg picture for time evolution. An object in a given Heisenberg state defines a four-dimensional reference frame and not a set of three-dimensional frames depending on time. In particular, the object defines the origin of the time scale with the precision permitted by the indeterminacy relations.

We indicate by $\mathcal{G}$ the Poincaré or the Galilei group, or some other space-time symmetry group. It acts on $\mathcal{H}^{(n)}$ by means of the unitary representation

$$g \rightarrow U^{(n)}(g) = U_0(g) \otimes U_1(g) \otimes \cdots \otimes U_{n-1}(g), \quad g \in \mathcal{G}. \tag{4}$$

The description of the system by means of the algebra $\mathcal{A}^{(n)}$ implicitly assumes the existence of an “external” object with a very large mass which defines a classical reference frame. If a classical reference frame is not available, only the $\mathcal{G}$ -invariant elements of $\mathcal{A}^{(n)}$ represent observables. In other words, $\mathcal{G}$ has to be considered as a global gauge group². We indicate by $\mathcal{B}^{(n)}$ the subalgebra of $\mathcal{A}^{(n)}$ composed of the $\mathcal{G}$ -invariant elements, namely the elements which commute with all the operators $U^{(n)}(g)$. It is a von Neumann algebra [17], namely it is symmetric (self-adjoint) and closed under the weak operator topology.

We also define the algebra $\mathcal{B}_k$ of the $\mathcal{G}$ -invariant elements of $\mathcal{A}_k$ which describes the “internal” degrees of freedom of the object F_k and the algebra $\mathcal{B}_{ik}$ of the invariant elements of $\mathcal{A}_{ik}$ which describes, besides the “internal” degrees of freedom of the objects F_i and F_k , the “relative” parameters, which define the relative position, time and velocity of the two objects. These relative parameters are the quantum analogs of the parameters defining the element of $\mathcal{G}$ which connects two classical reference frames. The homomorphisms $j_k^{(n)}$ and $j_{ik}^{(n)}$ can be restricted to these invariant subalgebras. We put

$$j_k^{(n)} \mathcal{B}_k = \mathcal{B}_k^{(n)} \subset \mathcal{B}^{(n)}, \quad j_{ik}^{(n)} \mathcal{B}_{ik} = \mathcal{B}_{ik}^{(n)} = \mathcal{B}_{ki}^{(n)} \subset \mathcal{B}^{(n)}. \tag{5}$$

²We do not consider a local Poincaré gauge invariance, which would lead to a theory of gravitation [16].

If i, j, k , are different indices, we have

$$\mathcal{B}_{ji}^{(n)} \cap \mathcal{B}_{jk}^{(n)} = \mathcal{B}_j^{(n)}, \quad n \geq 3. \quad (6)$$

Not all the kinds of objects are suitable for the definition of a reference frame. For instance, a spherically symmetric object cannot specify the directions of the spatial coordinate axes. If the object F_j defines a reference frame accurately, we expect that the algebra $\mathcal{B}_{jk}^{(n)}$ describes the object F_k completely and that the following property holds:

Assumption 1 *The algebra $\mathcal{B}^{(n)}$ is generated by the $n - 1$ subalgebras $\mathcal{B}_{jk}^{(n)}$ with j fixed and $k \neq j$.*

In the following Sections we shall test this assumption in some simple models. Without any loss of generality, we can assume that $j = 0$.

We remember [17] that the von Neumann algebra generated by a symmetric (self-adjoint) set $\mathcal{S}$ of operators is the double commutant $\mathcal{S}''$. If we indicate by $\mathcal{U}^{(n)}$ the von Neumann algebra generated by all the operators $U^{(n)}(g)$, we have

$$\mathcal{B}^{(n)} = \mathcal{U}^{(n)'}. \quad (7)$$

It is often convenient to describe a quantum system in terms of unbounded observables, which correspond to unbounded self-adjoint operators. If $\mathcal{S}$ is a set of self-adjoint not necessarily bounded operators, we indicate by $\mathcal{S}'$ the algebra of the bounded operators which commute with all the spectral projectors of all the operators in $\mathcal{S}$. We say that a self-adjoint operator is associated to a von Neumann algebra if its spectral projectors belong to the algebra.

If we indicate by $\mathcal{S}_{ik}$ a set of self-adjoint operators associated to $\mathcal{B}_{ik}^{(n)}$ and we have

$$\mathcal{S}'_0 = \mathcal{U}^{(n)}, \quad \mathcal{S}_0 = \bigcup_{k=1}^{n-1} \mathcal{S}_{0k}, \quad (8)$$

we obtain

$$\mathcal{S}''_0 = \mathcal{B}^{(n)} \quad (9)$$

and the Assumption 1 is satisfied for the frame F_0 . The same reasoning for $n = 2$ shows that $\mathcal{B}^{(2)} = \mathcal{B}_{01}$ is generated by $\mathcal{S}_{01}$ if $\mathcal{S}'_{01} = \mathcal{U}^{(2)}$.

We say that two subalgebras commute if all the elements of the first subalgebra commute with all the elements of the second one. One can easily see

that if the indices i, k, j, l are all different, the subalgebras $\mathcal{B}_{ik}^{(n)}$ and $\mathcal{B}_{jl}^{(n)}$ commute. However in a quantum theory one cannot assume that the subalgebras $\mathcal{B}_{jk}^{(n)}$ and $\mathcal{B}_{jl}^{(n)}$ commute. In fact we have:

Proposition 1 *If the objects F_0, F_1 and F_2 define quantum frames, namely they satisfy the Assumption 1, and all the pairs of the subalgebras $\mathcal{B}_{01}^{(3)}, \mathcal{B}_{02}^{(3)}$ and $\mathcal{B}_{12}^{(3)}$ commute, then the algebra $\mathcal{B}^{(3)}$ is commutative and describes a classical system.*

In fact, since the subalgebras $\mathcal{B}_{01}^{(3)}$ and $\mathcal{B}_{02}^{(3)}$ generate $\mathcal{B}^{(3)}$, $\mathcal{B}_{12}^{(3)}$ commutes with $\mathcal{B}^{(3)}$. In a similar way we see that $\mathcal{B}_{02}^{(3)}$ commutes with $\mathcal{B}^{(3)}$. A third application of the Assumption 1 shows that $\mathcal{B}^{(3)}$ commutes with itself, namely it is commutative.

Our aim is to study the relations between quantum frames and therefore we consider a set of n objects all of the same kind³ which describe n quantum frames and therefore satisfy the Assumption 1. We can simplify the notations by means of the identifications $\mathcal{B}_k = \mathcal{B}^{(1)}$ and $\mathcal{B}_{ik} = \mathcal{B}^{(2)}$. In order to obtain a compact notation for the various homomorphisms of the algebras $\mathcal{B}^{(n)}$, we introduce the sets $\Delta_n = \{0, \dots, n-1\}$ and for $n \geq m \geq 1$ the sets G_{nm} of the injective functions $\pi : \Delta_m \rightarrow \Delta_n$. If $\pi \in G_{nm}$, we consider the injective homomorphism $j_\pi : \mathcal{A}^{(m)} \rightarrow \mathcal{A}^{(n)}$ defined by

$$\begin{aligned} j_\pi(A_0 \otimes \dots \otimes A_{m-1}) &= B_0 \otimes \dots \otimes B_{n-1}, \\ B_{\pi(k)} &= A_k, \quad B_j = 1 \quad \text{for } j \notin \pi(\Delta_m). \end{aligned} \quad (10)$$

These homomorphisms can be restricted to the $\mathcal{G}$ -invariant subalgebras, namely we have $j_\pi : \mathcal{B}^{(m)} \rightarrow \mathcal{B}^{(n)}$. They have the property

$$j_\pi j_\sigma = j_{\pi \circ \sigma}, \quad \pi \in G_{np}, \quad \sigma \in G_{pm}, \quad n \geq p \geq m \geq 1. \quad (11)$$

We have

$$j_k^{(n)} = j_\pi, \quad \pi \in G_{n1}, \quad \pi(0) = k, \quad (12)$$

$$j_{ik}^{(n)} = j_\pi, \quad \pi \in G_{n2}, \quad \pi(0) = i, \quad \pi(1) = k. \quad (13)$$

We introduce the automorphism s of the algebra $\mathcal{B}^{(2)}$ by means of the formula

$$s = j_\pi, \quad \pi \in G_{2,2}, \quad \pi(0) = 1, \quad \pi(1) = 0. \quad (14)$$

³We assume, however, that they have some different quantum number in order to avoid the symmetrization or the antisymmetrization of the wave function.

and from eq. (11) we get

$$j_{ki}^{(n)} s = j_{ik}^{(n)}, \quad s^2 = 1. \quad (15)$$

The case $n = 3$ is particularly interesting: we have an algebra $\mathcal{B}^{(3)}$ that contains three isomorphic subalgebras $\mathcal{B}_{01}^{(3)}$, $\mathcal{B}_{02}^{(3)}$ and $\mathcal{B}_{12}^{(3)}$, and is generated by any pair of them. This means, for instance, that an element of $\mathcal{B}_{12}^{(3)}$, which concerns the relation between the frames F_1 and F_2 , can be expressed as a weak limit of algebraic expressions containing elements of $\mathcal{B}_{01}^{(3)}$ and $\mathcal{B}_{02}^{(3)}$. In other words, the mathematical structure of $\mathcal{B}^{(3)}$ with its three subalgebras describes the composition law of the transformations of quantum frames, which can be considered as the quantum version of the multiplication law of the group $\mathcal{G}$. It is not yet clear if the algebras $\mathcal{B}^{(n)}$ with $n > 3$ contain further physically relevant informations. One can remark that the formulation of the associative law requires three transformations and four reference frames.

In the treatment given above the algebras $\mathcal{B}^{(n)}$ and the homomorphisms j_π have been constructed starting from the group $\mathcal{G}$ and its unitary representations. One can conceive a more general kind of theory in which the group $\mathcal{G}$ does not exist and the relations between quantum reference frames are described directly by the von Neumann algebras $\mathcal{B}^{(n)}$ and by the homomorphisms j_π . Of course, some properties which follow from the construction based on the group $\mathcal{G}$ have to be assumed explicitly. Besides the Assumption 1, we require:

Assumption 2 *The homomorphisms j_π are injective and satisfy eq. (11).*

Assumption 3 *If the indices i, k, j, l are all different, the subalgebras $\mathcal{B}_{ik}^{(n)}$ and $\mathcal{B}_{jl}^{(n)}$ commute.*

One can also assume that eq. (6) is valid, possibly after a redefinition of the algebra $\mathcal{B}^{(1)}$. Note that our assumptions concern only the algebras, without any reference to the underlying Hilbert spaces.

A special case of eq. (11) is

$$j_\pi j_{ik}^{(m)} = j_{\pi(i), \pi(k)}^{(n)}, \quad \pi \in G_{nm}, \quad n \geq m \geq 2 \quad (16)$$

and from the Assumption 1, we see that the homomorphism j_π is uniquely determined by the homomorphism $j_{ik}^{(n)}$ and $j_{ik}^{(m)}$. Note however that a homomorphism j_π that satisfies eq. (16) does not necessarily exist.

From our assumptions one can derive the following result, that is stronger than Proposition 1:

Proposition 2 *If the indices i, k, j are all different and the subalgebras $\mathcal{B}_{jk}^{(n)}$ and $\mathcal{B}_{ji}^{(n)}$ commute, then all the algebras $\mathcal{B}^{(m)}$ with $m \leq n$ are commutative.*

The automorphisms j_π with $\pi \in G_{nn}$ permute the subalgebras $\mathcal{B}_{jk}^{(n)}$ and, taking into account the Assumption 3, we see that all the pairs of subalgebras $\mathcal{B}_{ik}^{(n)}$ commute. As in the proof of Proposition 1, it follows that all the subalgebras $\mathcal{B}_{jk}^{(n)}$ belong to the centre of $\mathcal{B}^{(n)}$ and that $\mathcal{B}^{(n)}$ is commutative. The algebras $\mathcal{B}^{(m)}$ with $m < n$ are isomorphic to subalgebras of $\mathcal{B}^{(n)}$ and the Proposition is proven.

Proposition 2 means that in a quantum theory the observables which describe a transformation between the frames F_j and F_i cannot be compatible with the observables which describe a transformation between the frames F_j and F_k . This fact is strictly related to the “paradox of quantum frames” discussed in ref. [2].

3 Classical reference frames.

By considering the classical limit of the formalism developed in the preceding section, we clarify the meaning of various concepts and we can understand why the relations between quantum frames cannot be described by a quantum group [13, 14] obtained by a “quantization” of the group $\mathcal{G}$.

In a classical theory the algebras $\mathcal{A}_0, \dots, \mathcal{A}_{n-1}$ and $\mathcal{A}^{(n)}$ are commutative and can be considered as algebras of L^∞ functions (essentially bounded measurable functions) on the phase spaces $\Gamma_0, \dots, \Gamma_{n-1}$ and

$$\Gamma^{(n)} = \Gamma_0 \times \Gamma_1 \times \dots \times \Gamma_{n-1}, \quad (17)$$

endowed with their Liouville measures. If we adopt the weak topology defined by the duality with the spaces L^1 , we are dealing with commutative von Neumann algebras of operators acting multiplicatively on the corresponding L^2 function spaces. The Lie group $\mathcal{G}$ acts smoothly on the phase spaces preserving the Liouville measures. For this action we use the notation $(g, x_k) \rightarrow gx_k$, $g \in \mathcal{G}$, $x_k \in \Gamma_k$. A function $f \in \mathcal{B}^{(n)}$ has the invariance property

$$f(gx_0, \dots, gx_{n-1}) = f(x_0, \dots, x_{n-1}). \quad (18)$$

We say that $\mathcal{G}$ acts freely on Γ_0 if, for any $x_0 \in \Gamma_0$, $gx_0 = x_0$ implies $g = e$ (e is the unit element). We also say that two submanifolds intersect transversely at a point if the vectors tangent to the submanifolds at this point generate the whole tangent space. Then we have:

Proposition 3 *If $\mathcal{G}$ acts freely on Γ_0 and there is a submanifold $\hat{\Gamma}_0 \subset \Gamma_0$ that intersects transversely every orbit of Γ_0 in one point, $\mathcal{B}^{(n)}$ is generated by the $n - 1$ subalgebras $\mathcal{B}_{0k}^{(n)}$, ($k = 1, \dots, n - 1$). Under these conditions the Assumption 1 is satisfied by the frame F_0 .*

This result could be obtained under much weaker hypotheses, but our purpose is only to give an example.

In order to sketch a proof, we remark that every $x_0 \in \Gamma_0$ can be written uniquely in the form $x_0 = gy_0$, with $y_0 \in \hat{\Gamma}_0$, and this relation defines a diffeomorphism of Γ_0 and $\mathcal{G} \times \hat{\Gamma}_0$. The Liouville measure on Γ_0 is transformed into the product of a left invariant measure on $\mathcal{G}$ and a suitable measure on $\hat{\Gamma}_0$. Since we have

$$f(gy_0, x_1, \dots, x_n) = f(y_0, g^{-1}x_1, \dots, g^{-1}x_{n-1}), \quad f \in \mathcal{B}^{(n)}, \quad (19)$$

we can identify the algebra $\mathcal{B}^{(n)}$ with the algebra of the L^∞ functions defined on $\hat{\Gamma}_0 \times \Gamma_1 \times \dots \times \Gamma_{n-1}$. The subalgebra generated by the subalgebras $\mathcal{B}_{0k}^{(n)}$ contains all the products of L^∞ functions of the kind $f(y_0, x_k)$ and their linear combinations. This set of functions is dense in $\mathcal{B}^{(n)}$.

We consider the special case in which all the objects are of the same kind and we put $\Gamma_0 = \dots = \Gamma_{n-1} = \Gamma$. We also assume that $\mathcal{G}$ acts freely and transitively on Γ . This means that the objects have no “internal” degree of freedom and $\mathcal{G}$ and Γ are diffeomorphic. If we choose a point y of Γ , and we put

$$f(g_0y, g_1y, \dots, g_{n-1}y) = \hat{f}(g_0^{-1}g_1, \dots, g_0^{-1}g_{n-1}), \quad (20)$$

we can consider $\mathcal{B}^{(n)}$ as a space of functions of $n - 1$ arguments in $\mathcal{G}$. In particular, $\mathcal{B}^{(2)}$ is a space of functions on $\mathcal{G}$ and we can write

$$\mathcal{B}^{(n)} = \mathcal{B}^{(2)} \otimes \dots \otimes \mathcal{B}^{(2)}, \quad (n - 1 \text{ factors}). \quad (21)$$

If we restrict our attention to continuous functions, $\mathcal{B}^{(2)}$ has a structure of Hopf algebra [13, 14]. If we want to work with L^∞ functions, we have to renounce to the existence of a counit. We have

$$[j_\pi f](x_0, x_1, \dots, x_{n-1}) = f(x_{\pi(0)}, x_{\pi(1)}, \dots, x_{\pi(m-1)}), \quad \pi \in G_{nm}, \quad (22)$$

$$[j_\pi \hat{f}](g_1, \dots, g_{n-1}) = \hat{f}(g_{\pi(0)}^{-1} g_{\pi(1)}, \dots, g_{\pi(0)}^{-1} g_{\pi(m-1)}), \quad (23)$$

where in the right hand side we have to put $g_0 = e$. We consider some particular cases:

$$[s \hat{f}](g) = \hat{f}(g^{-1}), \quad \hat{f} \in \mathcal{B}^{(2)}, \quad (24)$$

namely the automorphism s is the antipode of the Hopf algebra $\mathcal{B}^{(2)}$;

$$[j_{12}^{(3)} \hat{f}](g_1, g_2) = \hat{f}(g_1^{-1} g_2), \quad (25)$$

namely

$$j_{12}^{(3)} = (s \otimes 1)C, \quad (26)$$

where $C : \mathcal{B}^{(2)} \rightarrow \mathcal{B}^{(2)} \otimes \mathcal{B}^{(2)} = \mathcal{B}^{(3)}$ is the coproduct of the Hopf algebra. From the formula

$$[j_{0k}^{(n)} \hat{f}](g_1, \dots, g_{n-1}) = \hat{f}(g_k), \quad (27)$$

we obtain

$$j_{01}^{(3)} A = A \otimes 1, \quad j_{02}^{(3)} A = 1 \otimes A, \quad A \in \mathcal{B}^{(2)}. \quad (28)$$

The formulas (26) and (28) seem very natural in a formalism based on a Hopf algebra, but their generalization to non-commutative Hopf algebras (quantum groups) is not possible. In fact, we see from eq. (28) that the subalgebras $\mathcal{B}_{01}^{(3)}$ and $\mathcal{B}_{02}^{(3)}$ commute and it follows from the Proposition 2 that $\mathcal{B}^{(3)}$ and $\mathcal{B}^{(2)}$ must be commutative. Moreover, from eqs. (28) and (6) we have that $\mathcal{B}_0^{(3)}$ and therefore $\mathcal{B}^{(1)}$ contain only the multiples of the unit element. We see that a quantum group cannot describe the “internal” degrees of freedom of the quantum frames and the quantum aspects which are required by Proposition 2.

4 The simplest example.

As a first example of the general formalism, we consider the case in which $\mathcal{G}$ contains only the time translations and the objects that define the frames are essentially clocks. The algebra $\mathcal{B}^{(n)}$ is composed of the elements of $\mathcal{A}^{(n)}$ which commute with the Hamiltonian. Time measurements and clocks in quantum mechanics have been discussed by several authors [18, 19, 20, 21, 22, 23, 24, 25].

The simplest realization of a clock is a system with one degree of freedom. We consider a set of n clocks of this kind and we describe the clock F_k by means of the canonical variables q_k, p_k with $(\hbar = 1)$. We have

$$[q_i, p_k] = i\delta_{ik}, \quad (29)$$

$$H = \sum_k H_k, \quad H_k = 2^{-1}p_k^2. \quad (30)$$

This system can be reinterpreted as a single free particle in an n -dimensional space. The momenta p_k and the components of the “angular momentum”

$$L_{ik} = q_i p_k - q_k p_i \quad (31)$$

are conserved self-adjoint operators. They satisfy the commutation relations (29) and

$$[L_{ik}, p_j] = i\delta_{ij}p_k - i\delta_{kj}p_i, \quad (32)$$

$$[L_{ik}, L_{rs}] = i\delta_{ir}L_{ks} - i\delta_{is}L_{kr} - i\delta_{kr}L_{is} + i\delta_{ks}L_{ir}. \quad (33)$$

With the notation introduced in Sect. 2, we put $\mathcal{S}_{jk} = \{p_j, p_k, L_{jk}\}$. The set $\mathcal{S}_0$ contains all the momenta p_k and the quantities L_{0k} with $k = 1, \dots, n-1$, which generate the whole rotation group. It follows that the commutant $\mathcal{S}'_0$ is composed of the rotation-invariant bounded functions of the momenta p_k , namely the bounded functions of the Hamiltonian H , which form, in the present case, the algebra $\mathcal{U}^{(n)}$. We have seen that the condition (8) is satisfied and the Assumption 1 is valid.

The observable q_k can be interpreted as the running time directly read on the clock, while p_k is the rate of the clock. In the classical case, the true time is given by the ratio $T_k = q_k p_k^{-1}$. In the quantum case one can define in a suitable linear subspace of $\mathcal{H}$ the Hermitean operator

$$T_k = (2p_k)^{-1}q_k + q_k(2p_k)^{-1}, \quad (34)$$

which satisfies the commutation relation

$$i[H_k, T_k] = 1, \quad (35)$$

or, more exactly,

$$\exp(itH_k)T_k\exp(-itH_k) = T_k + t. \quad (36)$$

An argument due to Pauli [18] shows that under these conditions T_k cannot be self-adjoint. In fact, if it is self-adjoint one can prove that

$$\exp(iET_k)H_k\exp(-iET_k) = H_k - E \quad (37)$$

and this equation contradicts the fact that H_k has a spectrum bounded from below. Note that if an operator is not self-adjoint, it has no spectral representation, one cannot define bounded functions of it and it cannot represent an observable which has an arbitrarily small dispersion on a complete set of states.

The relative time of two clocks is given by

$$T_i - T_k = (2p_i p_k)^{-1} L_{ik} + L_{ik} (2p_i p_k)^{-1}. \quad (38)$$

From the commutation relations (32), we see that if we fix p_i or p_k , the quantity L_{ik} and the relative time are completely undetermined. We see that the “internal” variables p_i play an important role and cannot be eliminated by imposing that they take a fixed value. The same remark holds for the more complicated models described in the following Sections.

5 One-dimensional non-relativistic frames.

Now we assume that $\mathcal{G}$ is the Galilei group in one space dimension. An object which identifies a reference frame must contain at least two particles. We consider n objects composed of two free particles and we indicate by M_k the total mass and by μ_k the reduced mass of the object F_k . We adopt as canonical variables the centre of mass coordinate X_k , the total momentum P_k , the relative coordinate $\mu_k^{-1/2} q_k$ and the variable p_k conjugated to q_k . The Hamiltonian is

$$H = \sum_k H_k, \quad H_k = (2M_k)^{-1} P_k^2 + 2^{-1} p_k^2. \quad (39)$$

The canonical variables q_k and p_k have been scaled in order to eliminate the reduced mass μ_k . They describe a clock as in the preceding section.

The generators of the time and space translations are H and the total momentum

$$P = \sum_k P_k, \quad (40)$$

while the Galilei boosts are generated by

$$K = - \sum_k M_k X_k. \quad (41)$$

Note that

$$[H, P] = 0, \quad [H, K] = iP, \quad [P, K] = i \sum_k M_k. \quad (42)$$

The $\mathcal{G}$ -invariant variables are the internal variables p_k , the relative variables L_{ik} discussed in the preceding section, the relative velocities of the centres of mass

$$V_{ik} = M_k^{-1} P_k - M_i^{-1} P_i, \quad (43)$$

and the relative quantities

$$Y_{ik} = p_k(X_k - X_i) - q_k V_{ik}. \quad (44)$$

The quantity

$$(2p_k)^{-1} Y_{ik} + Y_{ik} (2p_k)^{-1} = X_k - X_i - T_k V_{ik} \quad (45)$$

can be interpreted as the space coordinate of the origin of the frame F_k measured with respect to the frame F_i .

We treat the Hilbert space $\mathcal{H}^{(n)}$ as a space of L^2 functions defined on the $2n$ -dimensional space $\mathcal{P}$ of the variables p_k and P_k (we use the same symbol for these variables and for the corresponding multiplication operators). The operators p_k and V_{ik} are clearly self-adjoint. The operators L_{ik} and Y_{ik} are generators of a group of linear measure-preserving transformations of the space $\mathcal{P}$ and they can be considered in a natural way as self-adjoint. One can see by means of a direct calculation that, if one excludes the lower dimensional manifold where $p_k = 0$ and $V_{ik} = 0$, the orbits of this group have codimension two and are just the manifolds on which the quantities H and P take constant values.

We put $\mathcal{S}_{ik} = \{p_i, p_k, L_{ik}, V_{ik}, Y_{ik}\}$ and we see that $\mathcal{S}_0$ contains all the “internal” variables p_k and the relative variables L_{0k} , V_{0k} and Y_{0k} . We replace in the space $\mathcal{P}$ the coordinates P_k by their linear combinations P and V_{0k} . If we indicate by $\mathcal{H}'$ the space of the L^2 functions of the variable P and by $\mathcal{H}''$ the space of the L^2 functions of the other variables q_k and V_{0k} , we have

$\mathcal{H}^{(n)} = \mathcal{H}' \otimes \mathcal{H}''$. An operator $A \in \mathcal{S}'_0$ is diagonal in the variables p_k and V_{0k} , but acts in a general way on the variable P . In other words, we have a direct integral decomposition [17] of A into operators $A(p_k, V_{0k}) \in \mathcal{W}$, where $\mathcal{W}$ is the algebra of the bounded operators acting on $\mathcal{H}'$. Since A must commute with L_{0k} and Y_{0k} , $A(p_k, V_{0k})$ must be constant on the orbits of the linear group generated by these operators, namely it depends only on the variable

$$\hat{H} = H - (2 \sum_k M_k)^{-1} P^2 = 2^{-1} \sum_k p_k^2 + (4 \sum_k M_k)^{-1} \sum_{ij} M_i M_j V_{ij}^2. \quad (46)$$

In conclusion, the elements of $\mathcal{S}'_0$ are given as direct integrals of bounded measurable functions $A(\hat{H})$ with values in $\mathcal{W}$ and we can write [17] $\mathcal{S}'_j = \mathcal{W} \otimes \mathcal{V}$, where $\mathcal{V}$ is the algebra of the bounded functions of $\hat{H}$ considered as operators on $\mathcal{H}''$. Note that the algebra $\mathcal{W}$ is generated by the operators P and

$$K = i \sum_k M_k \frac{\partial}{\partial P}. \quad (47)$$

It follows that $\mathcal{S}'_0$ is generated by the operators P , K and $\hat{H}$, namely it coincides with $\mathcal{U}^{(n)}$. Also in this case the condition (8) is satisfied and the Assumption 1 is valid.

6 One-dimensional relativistic frames.

As in the non relativistic case, we consider n frames, each one defined by means of two free particles, described by two irreducible unitary representations of the Poincaré group with masses m'_k and m''_k . The generators of the Poincaré transformations are given by

$$H = \sum_k H_k, \quad P = \sum_k P_k, \quad K = \sum_k K_k, \quad (48)$$

where (assuming $c = 1$)

$$\begin{aligned} H_k &= H'_k + H''_k = ((P'_k)^2 + (m'_k)^2)^{1/2} + ((P''_k)^2 + (m''_k)^2)^{1/2}, \\ P_k &= P'_k + P''_k, \quad K_k = K'_k + K''_k. \end{aligned} \quad (49)$$

The commutation relations are

$$[H, P] = 0, \quad [H, K] = iP, \quad [P, K] = iH \quad (50)$$

and similar relations hold for the generators acting on the various subsystems.

It is convenient to describe the object F_k by means of the variables H_k , P_k , K_k and

$$p_k = P'_k H''_k - P''_k H'_k, \quad q_k = H'_k K''_k - H''_k K'_k, \quad r_k = K'_k P''_k - K''_k P'_k, \quad (51)$$

which are connected by the identities

$$H_k^2 - P_k^2 = (m'_k)^2 + (m''_k)^2 + 2((m'_k m''_k)^2 + p_k^2)^{1/2} = M_k^2, \quad (52)$$

$$p_k K_k + q_k P_k + r_k H_k = 0. \quad (53)$$

We have the commutation relations

$$\begin{aligned} [H, p_k] &= 0, & [H, q_k] &= -ip_k, & [H, r_k] &= 0, \\ [P, p_k] &= 0, & [P, q_k] &= 0, & [P, r_k] &= -ip_k, \\ [K, p_k] &= 0, & [K, q_k] &= ir_k, & [K, r_k] &= iq_k. \end{aligned} \quad (54)$$

$$\begin{aligned} [p_k, q_k] &= -iH_k((m'_k m''_k)^2 + p_k^2)^{1/2}, \\ [r_k, p_k] &= -iP_k((m'_k m''_k)^2 + p_k^2)^{1/2}, \\ [q_k, r_k] &= -iK_k((m'_k m''_k)^2 + p_k^2)^{1/2}. \end{aligned} \quad (55)$$

These relations and the non-relativistic limit suggest the interpretation of the quantities

$$\hat{T}_k = t - T_k, \quad \hat{X}_k = (2p_k)^{-1} r_k + r_k (2p_k)^{-1}, \quad (56)$$

where T_k is given by eq. (34), as the time and space coordinates of the origin of the frame F_k measured in a classical frame. The relative coordinates of the frames F_k and F_i are given by

$$\begin{aligned} \hat{T}_k - \hat{T}_i &= (2p_i p_k)^{-1} Q_{ik} + Q_{ik} (2p_i p_k)^{-1}, \\ \hat{X}_k - \hat{X}_i &= (2p_i p_k)^{-1} R_{ik} + R_{ik} (2p_i p_k)^{-1}, \end{aligned} \quad (57)$$

where

$$Q_{ik} = q_i p_k - q_k p_i, \quad R_{ik} = p_i r_k - p_k r_i. \quad (58)$$

These quantities commute with H and P and satisfy the commutation relations

$$[K, Q_{ik}] = -iR_{ik}, \quad [K, R_{ik}] = -iQ_{ik}. \quad (59)$$

It follows that the quantities

$$L_{ik} = (H_i + H_k)Q_{ik} - (P_i + P_k)R_{ik}, \quad (60)$$

$$Y_{ik} = (H_i + H_k)R_{ik} - (P_i + P_k)Q_{ik} \quad (61)$$

are Poincaré invariant. They are proportional to the relative time and space coordinates of the frames F_i and F_k measured in the center of mass frame of these two objects. Note that the symbols which appear in the present section have not exactly the same meaning as in the preceding sections.

Another Poincaré invariant quantity is the relative rapidity of two frames which is given by

$$V_{ik} = \zeta_k - \zeta_i, \quad \zeta_i = \tanh^{-1} \frac{P_i}{H_i}. \quad (62)$$

As in the preceding section, we put $\mathcal{S}_{ik} = \{p_i, p_k, V_{ik}, L_{ik}, Y_{ik}\}$ and $\mathcal{S}_0$ contains the operators p_k , V_{0k} , L_{0k} and Y_{0k} . We describe the vectors of the Hilbert space $\mathcal{H}^{(n)}$ by means of wave functions in momentum space $\mathcal{P}$, square integrable with respect to the Lorentz invariant measure

$$d\mu = \prod_{k=0}^{n-1} \frac{dP'_k}{2H'_k} \frac{dP''_k}{2H''_k} = \prod_{k=0}^{n-1} 2^{-2} ((m'_k m''_k)^2 + p_k^2)^{-1/2} dp_k d\zeta_k. \quad (63)$$

Then p_k and V_{ik} are self-adjoint multiplication operators. Since we have

$$K'_k = -iH'_k \frac{\partial}{\partial P'_k}, \quad K''_k = -iH''_k \frac{\partial}{\partial P''_k}, \quad (64)$$

we see that L_{ik} and Y_{ik} are first order differential operators with smooth coefficients. They generate one-parameter groups of (non linear) measure-preserving diffeomorphism of the compact set in $\mathcal{P}$ defined by the inequality $H \leq C$. These diffeomorphisms can be extended to the whole manifold $\mathcal{P}$ and they define continuous groups of unitary operators in $\mathcal{H}^{(n)}$. It follows that the generators L_{ik} and Y_{ik} are self-adjoint.

A detailed analysis of the group of diffeomorphism generated by L_{0k} and Y_{0k} shows that, if we exclude the lower dimensional manifold where all the variables p_k vanish, the orbits have codimension two and are characterized by constant values of H and P . The arguments proceeds in close analogy with the non-relativistic case. We introduce in the space $\mathcal{P}$ the coordinates ζ_0 , p_k

and V_{0k} and we write $\mathcal{H}^{(n)} = \mathcal{H}' \otimes \mathcal{H}''$, where $\mathcal{H}'$ contains the L^2 functions of ζ_0 and $\mathcal{H}''$ contains the L^2 functions of the other coordinates. An operator $A \in \mathcal{S}'_0$ is diagonal in the variables p_k and V_{0k} and can be represented as a direct integral of a function with values in the algebra $\mathcal{W}$ of the bounded operators in $\mathcal{H}'$. This function is constant on the orbits and therefore depends only on the quantity

$$M^2 = H^2 - P^2 = \sum_{ik} M_i M_k \cosh V_{ik}. \quad (65)$$

We have $\mathcal{S}'_0 = \mathcal{W} \otimes \mathcal{V}$, where $\mathcal{V}$ contains the bounded functions of M^2 and $\mathcal{W}$ is generated by the operators ζ_0 and

$$K = i \frac{\partial}{\partial \zeta_0}. \quad (66)$$

Then $\mathcal{S}'_0$ is generated by the operators ζ_0 , K and M^2 , and it coincides with $\mathcal{U}^{(n)}$. The condition (8) is satisfied and the Assumption 1 is valid.

References

- [1] M. Toller: *An Operational Analysis of the Space-Time Structure*. Nuovo Cimento **B 40** (1977) 27.
- [2] Y. Aharonov and T. Kaufherr: *Quantum Frames of Reference*. Phys. Rev. **D 30** (1984) 368.
- [3] C. Rovelli: *What is Observable in Classical and Quantum Gravity?* Class. Quantum Grav. **8** (1991) 297.
- [4] C. Rovelli: *Quantum Reference Systems*. Class. Quantum Grav. **8** (1991) 317.
- [5] C. A. Mead: *Possible Connection between Gravitation and Fundamental Length*. Phys. Rev. **135 B** (1964) 849.
- [6] B. Ferretti: *On the Existence of a Minorant of the Indefiniteness for the Measurement of a Position*. Lett. Nuovo Cimento **40** (1984) 169.

- [7] L. J. Garay: *Quantum Gravity and Minimum Length*. Int. J. Mod. Phys. **A 10** (1995) 145.
- [8] S. Doplicher, K. Fredenhagen and J. E. Roberts: *The Quantum Structure of Spacetime at the Planck Scale and Quantum Fields*. Commun. Math. Phys. **172** (1995) 187.
- [9] S. Majid: *Hopf Algebras for Physics at the Planck Scale*. Class. Quantum Grav. **5** (1988) 1587.
- [10] S. Giller, P. Kosinski, M. Majewski, P. Maslanka and J. Kunz: *More about the q -Deformed Poincaré Algebra*. Phys. Lett. **B 286** (1992) 57.
- [11] J. Lukierski, A. Nowicki and H. Ruegg: *New Quantum Poincaré Algebra and κ -Deformed Field Theory*. Phys. Lett. **B 293** (1992) 344.
- [12] M. Maggiore: *Quantum Groups, Gravity, and the Generalized Uncertainty Principle*. Phys. Rev. **D 49** (1994) 5182.
- [13] E. Abe: *Hopf Algebras*. Cambridge University Press (1977).
- [14] N. Yu. Reshetikhin, L. A. Takhtadzhyan and L. D. Faddeev: *Quantization of Lie Groups and Lie Algebras*. Leningrad Math. J. **1** (1990) 193.
- [15] R. Giles: *Foundations for Quantum Mechanics*. Journ. Math. Phys. **11** (1970) 2139.
- [16] F. W. Hehl: *Four Lectures on the Poincaré Gauge Field Theory*. in “Cosmology and Gravitation”, P. G. Bergmann and V. De Sabbata editors, Plenum, New York, 1980, p. 5. (1980).
- [17] J. Dixmier: *Les algèbres d’opérateurs dans l’espace Hilbertien (algèbres de von Neumann)*. Gauthier-Villars, Paris (1969).
- [18] W. Pauli: *Die allgemeinen Prinzipien der Wellenmechanik*. Handbuch der Physik, edited by S. Flügge, vol. V/1, p. 60, Springer Verlag, Berlin, 1958.
- [19] H. L. Armstrong: *On a Space-Time Representation of Nonrelativistic Wave Mechanics*. Am. J. Phys. **22** (1957) 195.

- [20] H. Salecker and E. P. Wigner: *Quantum Limitations to the Measurement of Space-Time*. Phys. Rev. **109** (1958) 571.
- [21] F. Engelman and E. Fick: *Die Zeit in der Quantenmechanik*. Suppl. Nuovo Cimento **12** (1959) 63.
- [22] Y. Aharonov and D. Bohm: *Time in Quantum Theory and the Uncertainty Relation for Time and Energy*. Phys. Rev. **122** (1961) 1649.
- [23] D. M. Rosenbaum: *Super Hilbert Space and the Quantum Mechanical Time Operators*. Journ. Math. Phys. **10** (1969) 1127.
- [24] V. S. Olkhowsky, E. Recami and A. J. Gerasimchuk: *Time Operator in Quantum Mechanics. I: Nonrelativistic Case*. Nuovo Cimento **22** (1974) 263.
- [25] C. Rovelli: *Quantum Mechanics Without Time: a Model*. Phys. Rev. **D 42** (1990) 2638; Phys. Rev. **D 44** (1991) 1339.